
Education

Ankle Sprain Rehabilitation Exercises

As soon as you can tolerate pressure on the ball of your foot, begin stretching your ankle using the towel stretch. When this stretch is too easy, try the standing calf stretch and soleus stretch.

- Towel stretch: Sit on a hard surface with your injured leg stretched out in front of you. Loop a towel around the ball of your foot and pull the towel toward your body keeping your knee straight. Hold this position for 15 to 30 seconds then relax. Repeat 3 times.
- Standing calf stretch: Facing a wall, put your hands against the wall at about eye level. Keep the injured leg back, the uninjured leg forward, and the heel of your injured leg on the floor. Turn your injured foot slightly inward (as if you were pigeon-toed) as you slowly lean into the wall until you feel a stretch in the back of your calf. Hold for 15 to 30 seconds. Repeat 3 times. Do this exercise several times each day.
- Standing soleus stretch: Stand facing a wall with your hands at about chest level. With both knees slightly bent and the injured foot back, gently lean into the wall until you feel a stretch in your lower calf. Once again, angle the toes of your injured foot slightly inward and keep your heel down on the floor. Hold this for 15 to 30 seconds. Return to the starting position. Repeat 3 times.

You can do the next 5 exercises when your ankle swelling has stopped increasing.

- Ankle range of motion: Sitting or lying down with your legs straight and your knee toward the ceiling, move your ankle up and down, in and out, and in circles. Only move your ankle. Don't move your leg. Repeat 10 times in each direction. Push hard in all directions.
- Resisted dorsiflexion: Sit with your injured leg out straight and your foot facing a doorway. Tie a loop in one end of the tubing. Put your foot through the loop so that the tubing goes around the arch of your foot. Tie a knot in the other end of the tubing and shut the knot in the door. Move backward until there is tension in the tubing. Keeping your knee straight, pull your foot toward your body, stretching the tubing. Slowly return to the starting position. Do 3 sets of 10.
- Resisted plantar flexion: Sit with your leg outstretched and loop the middle section of the tubing around the ball of your foot. Hold the ends of the tubing in both hands. Gently press the ball of your foot down and point your toes, stretching the tubing. Return to the starting position. Do 3 sets of 10.
- Resisted inversion: Sit with your legs out straight and cross your uninjured leg over your injured ankle. Wrap the tubing around the ball of your injured foot and then loop it around your uninjured foot so that the tubing is anchored there at one end. Hold the other end of the tubing in your hand. Turn your injured foot inward and upward. This will stretch the tubing. Return to the starting position. Do 3 sets of 10.
- Resisted eversion: Sit with both legs stretched out in front of you, with your feet about a shoulder's width apart. Tie a loop in one end of the tubing. Put your injured foot through the loop so that the tubing goes around the arch of that foot and wraps around the outside of the uninjured foot. Hold onto the other end of the tubing with your hand to provide tension. Turn your injured foot up and out. Make sure you keep your uninjured foot still so that it will allow the tubing to stretch as you move your injured foot. Return to the starting position. Do 3 sets of 10.

You may do the rest of the exercises when you can stand on your injured ankle without pain.

- Heel raises: Balance yourself while standing behind a chair or counter. Raise your body up onto your toes and hold for 5 seconds. Then slowly lower yourself down. When this exercise becomes less painful, try lowering on the injured leg only. Repeat 10 times. Do 3 sets of 10.
- Step-up: Stand with the foot of your injured leg on a support (like a block of wood) 3 to 5 inches high. Keep your other foot flat on the floor. Shift your weight onto the injured leg and straighten the knee as the uninjured leg comes off the floor. Lower your uninjured leg to the floor slowly. Do 3 sets of 10.
- Static and dynamic balance exercises
 1. Place a chair next to your non-injured leg and stand upright. (This will provide you with balance if needed.) Stand on your injured foot. Try to raise the arch of your foot while keeping your toes on the floor. Try to maintain this position and balance on your injured side for 30 seconds. This exercise can be made more difficult by doing it on a piece of foam or a pillow, or with your eyes closed.
 2. Stand in the same position as above. Keep your foot in this position and reach forward in front of you

- with your injured side's hand, allowing your knee to bend. Repeat this 10 times while maintaining the arch height. This exercise can be made more difficult by reaching farther in front of you. Do 2 sets.
- Stand in the same position as above. While maintaining your arch height, reach the injured side's hand across your body toward the chair. The farther you reach, the more challenging the exercise. Do 2 sets of 10.
- Jump rope: Jump rope landing on both legs for 5 minutes, then on only the injured leg for 5 minutes.

Sports Medicine Advisor 2006.4; Copyright © 2006 McKesson Corporation and/or one of its subsidiaries. All Rights Reserved. Written by Tammy White, MS, PT, and Phyllis Clapis, PT, DHSc, OCS, for McKesson Provider Technologies. This content is reviewed periodically and is subject to change as new health information becomes available. The information is intended to inform and educate and is not a replacement for medical evaluation, advice, diagnosis or treatment by a healthcare professional.

