

## Education

### Snapping Hip Syndrome Rehabilitation Exercises

You may do all of these exercises right away.


- Quadriceps stretch: Stand an arm's length away from the wall, facing straight ahead. Brace yourself by keeping the hand on the uninjured side against the wall. With your other hand, grasp the ankle of the injured leg and pull your heel toward your buttocks. Don't arch or twist your back and keep your knees together. Hold this stretch for 15 to 30 seconds. Repeat 3 times.
- Hamstring stretch on wall: Lie on your back with your buttocks close to a doorway, and extend your legs straight out in front of you along the floor. Raise the injured leg and rest it against the wall next to the door frame. Your other leg should extend through the doorway. You should feel a stretch in the back of your thigh. Hold this position for 15 to 30 seconds. Repeat 3 times.
- Piriformis stretch: Lying on your back with both knees bent, rest the ankle of your injured leg over the knee of your uninjured leg. Grasp the thigh of your uninjured leg and pull that knee toward your chest. You will feel a stretch along the buttocks and possibly along the outside of your hip on the injured side. Hold this for 15 to 30 seconds. Repeat 3 times.
- Iliotibial band stretch (standing): Cross your uninjured leg in front of your injured leg and bend down and touch your toes. You can move your hands across the floor toward the uninjured side and you will feel more stretch on the outside of your thigh on the injured side. Hold this position for 15 to 30 seconds. Return to the starting position. Repeat 3 times.
- Iliotibial band stretch (side-leaning): Stand sideways near a wall, your injured leg toward the inside. Place the hand of your injured side on the wall for support. Cross your uninjured leg over the injured leg, keeping the foot of the injured leg stable. Lean into the wall. Hold the stretch for 15 seconds and repeat 3 times.
- Prone hip extension: Lie on your stomach with your legs straight out behind you. Tighten up your buttocks muscles and lift one leg off the floor about 8 inches. Keep your knee straight. Hold for 5 seconds. Then lower your leg and relax. Do 3 sets of 10.
- Side-lying leg lift: Lying on your uninjured side, tighten the front thigh muscles on your injured leg and lift that leg 8 to 10 inches away from the other leg. Keep the leg straight. Do 3 sets of 10.

*Sports Medicine Advisor 2006.4; Copyright © 2006 McKesson Corporation and/or one of its subsidiaries. All Rights Reserved. Written by Tammy White, MS, PT, and Phyllis Clapis, PT, DHSc, OCS, for McKesson Provider Technologies. This content is reviewed periodically and is subject to change as new health information becomes available. The information is intended to inform and educate and is not a replacement for medical evaluation, advice, diagnosis or treatment by a healthcare professional.*

## Snapping Hip Exercises



**Quadriceps stretch**



**Hamstring stretch on wall**



**Piriformis stretch**



**Standing**



**Side-leaning**

**Iliotibial band stretches**



**Prone hip extension**



**Side-lying leg lift**